

3. **SEBAEA** Solander ex R. Brown, Prodr. 451. 1810.

小黄管属 xiao huang guan shu

Annuals. Leaves opposite; leaf blade scalelike to lanceolate. Cymes terminal. Flowers 5-merous. Calyx lobed nearly to base. Corolla lobes shorter than tube. Stamens inserted at throat of corolla tube just below sinus between corolla lobes; anthers dehiscent longitudinally. Ovary 2-locular. Capsules 2-valved, many seeded. Seed coat alveolate.

About 100 species: temperate Africa, Bhutan, China, India, Madagascar, Nepal, Sri Lanka; one species in China.

1. **Sebaea microphylla** (Edgeworth) Knoblauch, Bot. Centralbl. 60: 324. 1894.

小黄管 xiao huang guan

Cicendia microphylla Edgeworth, Trans. Linn. Soc. London 20: 83. 1846; *Sebaea khasiana* C. B. Clarke.

Plants 5–10 cm tall, glabrous. Stems erect, simple, terete. Leaves sessile, 1–2.5 × 0.5–0.8 mm, base obtuse, margin membranous, apex acuminate. Flowers 1–3. Pedicel 0.7–1.7 cm, slender. Calyx 4–5 mm; lobes spreading, lanceolate, 3.5–4.5 mm, margin broadly membranous, apex acuminate, midvein keeled. Corolla yellow, tubular, 8–10 mm; lobes elliptic, 4–5 mm, margin entire, apex obtuse. Filaments linear, 1.8–2 mm; anthers ellipsoid, 0.8–1 mm. Style linear, 3.5–4 mm; stigma lobes orbicular. Capsules ellipsoid to subglobose, 2.5–3 mm. Seeds dark brown, ellipsoid to globose, 0.2–0.3 mm. Fl. and fr. Oct.

Forests; ca. 2300 m. Yunnan [Bhutan, India, Nepal].

